

Quarkonia Production in Heavy Ion Collisions

Che-Ming Ko,
Texas A&M University

- In-medium properties of quarkonia
- Quarkonia production mechanisms in HIC
- Nuclear modification factor for J/ψ
- Nuclear modification factor for $Y(1S)$
- J/ψ elliptic flow
- Effects of initial fluctuation on bottomonia production
- Hot medium effects in p+Pb

In collaboration with Taesoo Song, Yunpeng Liu, and Kyongchol Han

Support by US National Science Foundation and Department of Energy, and
The Welch Foundation

Quarkonia in QGP

Free energy F for a pair of $Q\bar{Q}$ from LQCD
 [Kaczmarek, EJP 61, 811 (2009)]

Two limits of the potential:

$$V(r, T) = F$$

$$\text{or } V(r, T) = U = F + TS$$




Schroedinger equation at finite T :

binding energy $\epsilon(T)$
 radius $R(T)$

Dissociation temperature:

$$\epsilon(T_D) \rightarrow 0, R(T_D) \rightarrow \infty$$

For $V=U$ (Satz et al.)





state	$J/\psi(1S)$	$\chi_c(1P)$	$\psi'(2S)$	$\Upsilon(1S)$	$\chi_b(1P)$	$\Upsilon(2S)$	$\chi_b(2P)$	$\Upsilon(3S)$
T_d/T_c	2.10	1.16	1.12	> 4.0	1.76	1.60	1.19	1.17

QCD sum rule study of J/ψ

Lee, Morita, Song & Ko,
PRD 89, 094015 (2014)

$$\Pi_{\mu\nu}(q) = i \int d^4x e^{iqx} \langle T[\bar{c}(x)\gamma_\mu c(x)\bar{c}(0)\gamma_\nu c(0)] \rangle.$$



- Results favor free energy as the potential between charm and anticharm quarks

Screened Cornell potential for heavy quark and antiquark in QGP


- Screened Cornell potential between charm and anticharm quarks

$$V(r,T) = \frac{\sigma}{\mu(T)} \left[1 - e^{-\mu(T)r} \right] - \frac{\alpha}{r} e^{-\mu(T)r}$$

with string tension $\sigma = 0.192 \text{ GeV}^2$
and screening mass

$$\mu(T) = \sqrt{\frac{N_c}{3} + \frac{N_f}{6}} gT$$

- Its strength is between the internal energy (U) and free energy (F) of heavy quark and antiquark from LQCD; similar to F at T_c and to U at $4T_c$.



Thermal properties of charmonia

- Binding energy



$$\varepsilon_0 = 2m_c + \frac{\sigma}{\mu(T)} - E$$

Charm quark mass $m_c=1.32$ GeV


E: eigenvalues of Cornell potential

- Dissociation temperature T_D :
corresponding to $\varepsilon_0=0$

For $g=1.87$, $T_D \sim 300$ MeV for J/ψ
and $\sim T_D=175$ MeV for ψ' and χ_c




Thermal properties of bottomonia



State	$Y(1S)$	$\chi_b(1P)$	$Y'(2S)$	$\chi_b'(2P)$	$Y''(3S)$
Dissociation temp (T_c)	4	1.51	1.67	1.09	1.12

Quasiparticle model for QGP

P. Levai and U. Heinz, PRC , 1879 (1998)



$$p(T) = \sum_{i=g,q,\bar{q}} \frac{g_i}{6\pi^2} \int_0^\infty dk f_i(T) \frac{k^4}{E_i} - B(T)$$

$$e(T) = \sum_{i=g,q,\bar{q}} \frac{g_i}{2\pi^2} \int_0^\infty dk k^2 f_i(T) E_i + B(T)$$

$$m_g^2 = \left(\frac{N_c}{3} + \frac{N_f}{6} \right) \frac{g^2(T) T^2}{2}$$

$$m_q^2 = \frac{g^2(T) T^2}{3}$$

$$g^2(T) = \frac{48\pi^2}{(11N_c - 2N_f) \ln F^2(T, T_c, \Lambda)}$$


$$F(T, T_c, \Lambda) = \frac{18}{18.4 e^{-(T/T_c)^2/2} + 1} \frac{T}{T_c} \frac{T_c}{\Lambda}$$

- Resulting EOS is similar to that from LQCD by the hot QCD collaboration, and the difference is smaller than that between the hot QCD and Wuppertal-Budapest Collaborations

Thermal decay widths of quarkonia

Song, Park & Lee, PRC 81, 034914 (10)


- Dissociation by partons (NLO pQCD)



$$|\overline{M}|^2 = \frac{4}{3} g^4 m_c^2 m_{J/\psi} \left| \frac{\partial \psi(p)}{\partial p} \right|^2 \left\{ -\frac{1}{2} + \frac{(k_1^0)^2 + (k_2^0)^2}{2k_1 \cdot k_2} \right\}$$



- Dissociation by hadrons

$$\sigma(s) = \sum_i \int dx n_i(x, Q^2) \sigma_i(xs, Q^2)$$




- Thermal dissociation width

$$\Gamma(T) = \sum_i \int \frac{d^3k}{(2\pi)^3} v_{rel}(k) n_i(k, T) \sigma_i^{diss}(k, T)$$



Validity of dipole approximation in J/ ψ dissociation

Liu, Ko & Song, PRC 88, 046902 (2013)



- Dipole approximation overestimates J/ ψ dissociation cross section by as large as a factor of three

Directly produced J/ψ

Song, Park & Lee,
PRC 81, 034914 (10)

- Number of initially produced

$$N_{J/\psi}^{AA} = \sigma_{J/\psi}^{NN} A^2 T_{AA}(\vec{b})$$

- $\sigma_{J/\psi}^{NN}$: J/ψ production cross section in NN collision; $\sim 0.774 \mu\text{b}$ at $s^{1/2} = 200 \text{ GeV}$

- Overlap function

$$T_{AA}(\vec{b}) = \int d^2\vec{s} T_A(\vec{s}) T_A(\vec{b} - \vec{s})$$

- Thickness function

$$T_A(\vec{s}) = \int_{-\infty}^{\infty} dz \rho_A(\vec{s}, z)$$

- Normalized density distribution

$$\rho(r) = \frac{\rho_0}{1 + e^{(r-r_0)/c}}$$

$r_0 = 6.38 \text{ fm}$, $c = 0.535 \text{ fm}$ for Au


- Nuclear absorption

- Survival probability

$$S_{nuct}(\vec{b}, \vec{s}) = \frac{1}{T_{AB}(\vec{b})} \int dz dz' \rho_A(\vec{s}, z) \rho_B(\vec{b} - \vec{s}, z')$$

$$\times \exp \left\{ -(A-1) \int_z^{\infty} dz_A \rho_A(\vec{s}, z_A) \sigma_{nuc} \right\}$$

$$\times \exp \left\{ -(B-1) \int_{z'}^{\infty} dz_B \rho_B(\vec{s}, z_B) \sigma_{nuc} \right\}$$



Regenerated J/ψ

Rate equation for J/ψ production $\frac{dN_i}{d\tau} = -\Gamma_i(N_i - N_i^{\text{eq}}), \quad N_i^{\text{eq}} = \gamma^2 R n_i^{\text{GC}} V$

▪ Charm fugacity is determined by

$$N_{c\bar{c}}^{AA} = \left[\frac{1}{2} \gamma m_o \frac{I_1(\gamma n_o V)}{I_0(\gamma n_o V)} + \gamma^2 n_h \right] V = \sigma_{c\bar{c}}^{NN} A^2 T_{AA}(\vec{b})$$



• $\sigma_{c\bar{c}}^{NN}$: charm production cross section in NN collision; $\sim 63.7 \mu\text{b}$ at $s^{1/2} = 200 \text{ GeV}$

▪ Charm relaxation factor

$$R = 1 - \exp \left\{ - \int_{\tau_0}^{\tau_{QGP}} d\tau \Gamma_c(T(\tau)) \right\}$$

$$\Gamma(T) = \sum_i \int \frac{d^3 k}{(2\pi)^3} v_{\text{rel}}(k) n_i(k, T) \times \sigma_i^{\text{diss}}(k, T) (1 - \vec{p} \cdot \vec{p}' / p^2)$$


as J/ψ is more likely to be formed if charm quarks are in thermal equilibrium



Approximately reproduced by non-equilibrium charm quarks from parton cascade [PRC 85, 954905 (12)]

Nonequilibrium effects on J/Ψ production


Song, Han & Ko, PRC 85, 054905 (2012)



- Nonequilibrium effects can be approximated by the relaxation factor $t/t_{\text{relaxation}}$

Nuclear modification factor for J/ψ

Song, Han & Ko, PRC 84, 034907 (2011)





	SPS	RHIC	LHC	LHC
				$p_T > 6.5$ GeV
production (μb)				
$d\sigma_{J/\psi}^{pp}/dy$	0.05	0.774	4.0	
$d\sigma_{c\bar{c}}^{pp}/dy$	5.7	119	615	
feed-down (%)				
f_{χ_c}	25	32	26.4	23.5
$f_{\psi'(2S)}$	8	9.6	5.6	5
f_b			11	21
nuclear absorp.				
σ_{abs} (mb)	4.18	2.8	0 or 2.8	

- Most J/ψ are survivors from initially produced
- Kink in R_{AA} is due to the onset of initial temperature above the J/ψ dissociation temperature in QGP
- Inclusion of shadowing reduces slightly R_{AA}

Nuclear modification factor for $\Upsilon(1S)$


Song, Han & Ko, PRC 85, 014902 (2012)




- Regeneration contribution is negligible
- Primordial excited bottomonia are largely dissociated
- Medium effects on bottomonia reduce R_{AA} of $\Upsilon(1S)$

Y(1S) nuclear modification factor at LHC from other models


1) Strickland, PRL 107, 132301 (2011)



2) Zhuang et al.,




3) Emerick, Zhao & Rapp, EPJA 48, 72 (2012)



- | | | |
|--------------------------------|------------------------------|-----------------------------|
| ▪ Potential: in-medium Cornell | ▪ Potential: U or F | ▪ Potential: ~ U or vacuum |
| ▪ Disso.: LO pQCD | ▪ Disso.: vacuum gluo-disso. | ▪ Diss.: vacuum gluo-disso. |
| ▪ Dynamics: anisotropic hydro | ▪ Dynamics: ideal hydro | ▪ Dynamics: fireball |

4) Brezinzki & Wolschin, PLB 707, 534 (12): estimate using in-medium gluo-dissociation

Thermal decay width of $Y(1S)$ in different models




- Thermal decay width
 - Rapp: quasielastic scattering
 - Zhuang: OPE by Peskin
 - Strickland: LO pQCD
 - Song: NLO pQCD

- Very different thermal decay widths are used in different models

J/ψ elliptic flow

Song, Lee, Xu & Ko, PRC 83, 014914 (11)



$$v_2 = \frac{\int d\varphi \cos(2\varphi) (dN / dy d^2 p_T)}{\int d\varphi (dN / dy d^2 p_T)}$$

$$= \frac{\int dA_T \cos(2\varphi) I_2(p_T \sinh \rho / T) K_1(m_T \cosh \rho / T)}{\int dA_T I_0(p_T \sinh \rho / T) K_1(m_T \cosh \rho / T)}$$

$\rho = \tanh(v_T) = \text{transverse rapidity}$

Introducing viscous effect at freeze out
 $T = 125 \text{ MeV}$

$$\Delta v = (v_x - v_y) \exp[-C p_T / n]$$




with $C = 1.14 \text{ GeV}^{-1}$ and $n = \text{number of quarks in a hadron}$

- Initially produced J/ψ have essentially vanishing v_2
- Regenerated J/ψ have large v_2
- Final J/ψ v_2 is small as most are initially produced

Effects of initial fluctuations on bottomonia production


Song, Han & Ko, NPA 897, 141 (2012)

2+1 ideal hydro




- Initial fluctuations affect R_{AA} of bottomonia in peripheral collisions and at low p_T

J/ψ production in p+Pb at $\sqrt{s_{NN}} = 5.02$ TeV



Liu, Song & Ko, PLB 728, 437 (2013)



- Most central 10% collisions from AMPT
- Including hot medium effects better describes data

Summary

- J/ψ survives up to $1.7 T_c$ and $Y(1S)$ survives up to $4 T_c$
- Most observed J/ψ and $Y(1S)$ are from primordially produced; contribution from regeneration is small at present HIC
- Various models with different assumptions can describe experimental data
- Elliptic flow of regenerated J/ψ is large, while that of directly produced ones is essentially zero. Studying v_2 of J/ψ is useful for distinguishing the mechanism for J/ψ production in HIC
- Initial fluctuations affect R_{AA} of bottomonia in peripheral collisions and at low p_T
- Hot medium effects describe better J/ψ data from p+Pb collisions